

Dynamics of a tunneling magnetic impurity: Kondo effect induced incoherence

L. Borda,¹ G. Zaránd,^{1,2}

¹Research Group of the Hungarian Academy of Sciences, Institute of Physics, TU Budapest, H-1521 Hungary,

²Lyman Physics laboratory, Harvard University, Cambridge MA 02145

(February 1, 2008)

We study how the formation of the Kondo compensation cloud influences the dynamical properties of a magnetic impurity that tunnels between two positions in a metal. The Kondo effect dynamically generates a strong tunneling impurity-conduction electron coupling, changes the temperature dependence of the tunneling rate, and may ultimately result in the destruction of the coherent motion of the particle at zero temperature. We find an interesting two-channel Kondo fixed point as well for a vanishing overlap between the electronic states that screen the magnetic impurity. We propose a number of systems where the predicted features could be observed.

75.20Hr, 72.15Qm, 71.10Hf

Tunneling of a heavy particle or some collective degree of freedom in a dissipative environment has been the subject of intense theoretical and experimental research in the past and is by now reasonably well-understood [1–7]. Maybe the most intriguing case is that of *ohmic* dissipation, where the particle usually couples to electron-hole excitations of a metallic environment. In this case the bare tunneling amplitude Δ_0 of the particle is strongly renormalized due to the dissipative environment, and becomes temperature dependent. In the simplest scenario, where the tunneling occurs between two sites, the effective tunneling displays a power-law behavior over a wide range of temperatures, $\Delta(T) \sim \Delta_0(T/\omega_0)^\alpha$, with ω_0 a high-energy cut-off of the order of the Debye frequency, and α a dimensionless coupling constant [1,2]. The renormalization of the tunneling amplitude is a consequence of Anderson's orthogonality catastrophe [8]: At any position the presence of the particle generates a screening cloud that consists of an *infinite* number of electron-hole excitations. The formation of this huge 'electronic polaron-cloud' slows down the particle, increases its mass, and thus decreases its tunneling amplitude. Depending on the specific value of the coupling α , the dynamics of the particle can be of three different kind: (a) If the coupling is small the particle moves with slightly damped *coherent* oscillations between the two positions at $T = 0$. (b) For $1/2 < \alpha < 1$ the motion of the particle becomes *incoherent*, while for even larger couplings (c) the particle becomes localized and cannot move from one well to the other.

At low temperatures and larger values of α , more complicated dynamical processes involving simultaneous tunneling and electron scattering (electron assisted tunneling) may become more relevant and can, in principle, delocalize the particle or lead to other types of strongly correlated states [9,10]. In this paper, however, we focus on the rather generic case of 'slow' TLS's [9], where the energy scale generated by these processes is extremely small, and they can therefore be neglected.

In the present paper we focus our attention to the

very interesting but poorly understood case of a tunneling *magnetic impurity* coupled to an ohmic environment [11,12]. Possible examples of such a system include a magnetic impurity tunneling between an STM tip and a metallic surface [13], a Kondo impurity in an amorphous region [14], a spin 1/2 quantum soliton interacting with a metallic environment [15], or charge tunneling in double quantum-dot systems [16,17]. In all these cases the spin of a magnetic impurity couples to the conduction electrons through an exchange interaction. This exchange interaction becomes renormalized as the temperature is lowered and may lead to the dynamical formation of a Kondo compensation cloud. Since the tunneling particle has to drag this cloud with itself, the exchange coupling will lead to a dynamically generated temperature dependent renormalization of the tunneling amplitude. In the present paper we study this interplay between the Kondo effect and the orbital motion of the TLS in detail.

For the sake of simplicity we focus our attention to the simplest possible case of a TLS, where tunneling takes place between two positions only, $\mathbf{R}_\pm$. Furthermore, though we also discuss the role of asymmetry to some extent, we mostly focus on spatially symmetrical TLS's. We show, in particular, that the Kondo effect associated with the magnetic degrees of freedom leads to a strong *temperature-dependence* of the exponent α , and may eventually induce a decoherent state. Destroying the Kondo cloud with a magnetic field one could then be able to drive the particle back to the coherent regime. The orbital motion, on the other hand, may lead to the appearance of a *two-channel Kondo* state under special conditions, where the impurity tunnels very fast back and forth and forms a Kondo state with the conduction electrons at *both* positions.

We describe the TLS by the tunneling Hamiltonian

$$H_{\text{tun}} = -(\Delta_0 T^x + \Delta_z T^z), \quad (1)$$

where the two pseudospin states $T^z = \pm 1/2$ correspond to the two tunneling positions, Δ_0 is the tunneling matrix element, and Δ_z describes the asymmetry of the TLS.

As a first approximation, we consider the exchange interaction between the tunneling particle and the electrons as completely local:

$$H_{\text{int}} = \sum_{q=\pm} J_q P_q \vec{S} (\Psi_q^\dagger \vec{\sigma} \Psi_q). \quad (2)$$

Here $P_\pm = 1/2(1 \pm 2T^z)$ projects out the TLS states at positions $\mathbf{R}_\pm$, $J_\pm$ is the exchange coupling at these positions, $\vec{S}$ denotes the spin operator of the impurity, and $\vec{\sigma}$ stands for the Pauli matrices. The field operators, $\Psi_{\pm,\mu}^\dagger = \int e^{i\mathbf{k}\mathbf{R}_\pm} c_{\mathbf{k}\mu}^\dagger d^3\mathbf{k}/(2\pi)^3$, create conduction electrons at $\mathbf{R}_\pm$ with spin μ . Including more complicated non-local interactions in Eq. (2) such as spin-assisted tunneling has not changed our results [11,16].

First we consider the case of a symmetric TLS with $J_- = J_+ = J$ and $\Delta_z = 0$. Following the standard procedure [10,18], the relevant conduction electron degrees of freedom can be represented simply by one-dimensional fermion fields [2], $c_{p\alpha\mu}$, obeying canonical anticommutation relations, $\{c_{p\alpha\mu}, c_{p'\alpha'\mu'}^\dagger\} = 2\pi \delta(p-p')\delta_{\alpha\alpha'}\delta_{\mu\mu'}$, and described by

$$H_{\text{el}} = \sum_{\alpha=e,o} \sum_{\mu=\uparrow,\downarrow} \int_{-k_F}^{k_F} \frac{dp}{2\pi} v_F p c_{p\alpha\mu}^\dagger c_{p\alpha\mu}. \quad (3)$$

In Eq. (3) the radial momentum p is measured from the Fermi momentum k_F , v_F is the Fermi velocity, $\alpha = \{e, o\}$ is the parity, and μ denotes the spin. Introducing the fields $\Psi_{\alpha,\mu} \equiv \int_{-k_F}^{k_F} c_{p\alpha\mu} dp/\sqrt{2\pi}$, Eq. (2) becomes

$$H_{\text{int}} = \frac{g}{2} \vec{S}(1+F) \Psi_e^\dagger \vec{\sigma} \Psi_e + \frac{g}{2} \vec{S}(1-F) \Psi_o^\dagger \vec{\sigma} \Psi_o + g \vec{S} \sqrt{1-F^2} T^z (\Psi_e^\dagger \vec{\sigma} \Psi_o + \Psi_o^\dagger \vec{\sigma} \Psi_e). \quad (4)$$

Here $F = \sin(k_F d)/k_F d$ measures the overlap of the states $\Psi_\pm$, with $d = |\mathbf{R}_+ - \mathbf{R}_-|$ the tunneling distance, and $g = Jk_F^2/2\pi^2$. For $d = 0$ this Hamiltonian obviously reduces to the single-channel Kondo model. In the following we set $v_F = k_F = \hbar = 1$.

To study the zero temperature dynamical properties of the tunneling Kondo spin we used Wilson's numerical renormalization group (NRG) [19]. In this very accurate numerical technique one constructs a series of Hamiltonians, H_N , which are diagonalized iteratively. Having obtained the many-body eigenstates and energies of H_N one can use them to calculate physical quantities at an energy scale $T, \omega \sim \omega_N \sim \Lambda^{-(N+1)/2}$, with $\Lambda \approx 3$ a discretization parameter. Our results were obtained by keeping the lowest 250 states in each iteration. To obtain accurate results we exploited the following symmetries: (i) parity (ii) global spin rotations, and (iii) a hidden $SU(2)$ symmetry, related to electron-hole symmetry [20].

We computed the $T = 0$ impurity spin and pseudospin spectral functions, $\varrho_{\mathcal{O}}^i(\omega) = -\text{Im}\{\chi_{\mathcal{O}}^i(\omega)\}$, (with $\chi_{\mathcal{O}}^i(\omega)$ the Fourier transform of the retarded response function), using their Lehmann representations:

$$\varrho_{\mathcal{O}}^i(\omega > 0) = \sum_n |\langle n | \mathcal{O}^i | 0 \rangle|^2 \delta(E_n - E_0 - \omega).$$

Here $|0\rangle$ ($|n\rangle$) is the ground (n th excited) state of the system with energy E_0 (E_n), and $\mathcal{O}^i$ denotes the operators T^i and S^i .




FIG. 1. Logarithmic plot of various spectral functions discussed in the text for $\Lambda = 3$, $g = 0.144$, and $\Delta_0 = 2.31 \cdot 10^{-5}$. The energy scales T_K and Δ are also indicated.

Our results are summarized in Fig. 1. Let us first focus on the somewhat peculiar case of $F = 0$, where Ψ_+ and Ψ_- do not overlap. In this case we can observe two distinct cross-overs: The first takes place at the Kondo energy $T_K \approx e^{-1/2g}$ and corresponds to the formation of a local Kondo state at the *actual* position of the TLS. Above T_K all spectral functions behave as $1/\omega$, indicating that all correlation functions are constant for times shorter than $1/T_K$. Below T_K the spin spectral function becomes linear. Performing a Hilbert transform one can see that this corresponds to a constant impurity susceptibility $\sim 1/T_K$. The z -component of the pseudospin spectral function is practically unaffected by the formation of the Kondo compensation cloud: at this time scale tunneling events are very rare, and the particle can be considered as immobile. This interpretation is confirmed by a detailed analysis of the finite size spectrum [16]. The logarithmic slope of the x -component of the pseudospin spectral function, however, does become modified: this change is related to the dynamical renormalization of the tunneling amplitude by the formation of the Kondo compensation cloud.

Though $\Delta_0 \ll T_K$ in our computations, Δ_0 is a relevant perturbation, and leads to a second cross-over at a renormalized value of the tunneling amplitude, Δ^* , where the TLS freezes into the even tunneling state. For $F = 0$ a two-channel Kondo state is formed below Δ^* , as confirmed by the analysis of the finite size spectrum [16].

This is most easily understood by observing that the last term of Eq. (4) flips the TLS between the $T^x = \pm 1/2$ states, and can therefore be dropped below Δ^* . Here H_{int} becomes simply the two-channel Kondo Hamiltonian, and a two-channel Kondo state is formed in the spin sector [9]. The spin spectral function and the z -component of the pseudospin spectral function become constant below Δ^* , implying the logarithmic divergence of the spin- and T^z -susceptibility at low temperatures, $\chi_S(T) \sim \Delta^* \ln(T_K/T)/T_K^2$, and $\chi_T^z(T) \sim \ln(T_K/T)/\Delta^*$, and the asymptotic behavior of the time-dependent correlation functions $\langle S^i(t)S^i(0) \rangle \sim \langle T^z(t)T^z(0) \rangle \sim 1/t$.

These results immediately imply that the external magnetic field and asymmetry are both relevant operators of scaling dimension $1/2$ at this two-channel Kondo fixed point [21]. The two-channel Kondo state is thus extremely unstable. It crosses over to a Fermi liquid state also for any finite overlap F : For small values of F a third crossover occurs at an energy T^* well below Δ^* , however, for generic F 's this second crossover takes place almost simultaneously with the crossover at Δ^* and only a small kink remains from the two-channel Kondo behavior of $F = 0$.

To determine the scale Δ^* and the temperature dependence of the tunneling rate we invoked renormalization group arguments. The Hamiltonian involves three dimensionless parameters: g , F , and Δ_0/ω_0 , with ω_0 the high-energy cutoff. Well above Δ^* , Δ_0/ω_0 is small and the effective tunneling at energy scale ω or temperature $T \sim \omega$, $\Delta(\omega)$, satisfies the following scaling equation:

$$\frac{d \ln \Delta(\omega)}{d \ln(\omega_0/\omega)} = -\alpha(g(\omega), F), \quad (\omega_0 \gg \Delta(\omega)). \quad (5)$$

Above Δ^* the impurity is immobile. Therefore the effective Kondo coupling $g(\omega)$ does not depend on the other two parameters, and is a universal function of ω/T_K . In this regime F can be completely transformed out of the Hamiltonian and is therefore constant. Thus $\alpha = \alpha(\omega/T_K, F)$ is a universal function of ω/T_K and F for $\omega_0 \gg \Delta$. This function is related to the time-dependent correlation function as $\langle T^x(t)T^x(0) \rangle \sim 1/t^{2\alpha(\omega)}$ ($t \sim 1/\omega$) [21]. Assuming that the retarded and the time ordered correlation functions have the same singular behavior, this immediately implies that, within logarithmic accuracy, α can be determined from the logarithmic derivative of ϱ_T^x as

$$\alpha(\omega) \approx \frac{1}{2} \left(\frac{d \ln \varrho_T^x}{d \ln \omega} + 1 \right), \quad (\omega \gg \Delta^*). \quad (6)$$

We plotted this universal function in Fig. 2. For large frequencies $\alpha \approx 0$, meaning that the tunneling amplitude remains unrenormalized above T_K . Below T_K , on the other hand, it scales to an overlap-dependent constant, $\alpha_<(F)$. As shown in the inset, $\alpha_<$ coincides with the Anderson orthogonality exponent K for a maximally strong scatterer with a phase shift $\delta_\uparrow = \delta_\downarrow = \pi/2$ [22],

$$\alpha_< \equiv \sum_{\sigma} \left(\frac{1}{\pi} \operatorname{atan} \frac{\tan \delta_{\sigma} \sqrt{1-F^2}}{\sqrt{1+F^2 \tan^2 \delta_{\sigma}}} \right)^2. \quad (7)$$

Thus, far below T_K (but still above Δ^*) the Kondo impurity acts as a maximally strong potential scatterer in agreement with Nozières' Fermi liquid picture [23].




FIG. 2. Top: Energy- and overlap-dependence of the anomalous dimension α of the dimensionless tunneling amplitude for $F = 0, 0.1, 0.2, 0.3, 0.4, 0.5$, and 0.6 (top to bottom), as determined from the logarithmic derivative of the spectral function ϱ_T^x . Bottom: Temperature dependence of the normalized tunneling rate, $\tilde{R}(T) = R(T)/\Delta_0^2$. F decreases from top to bottom. We used $\Lambda = 3$, $g = 0.144$, and $\Delta_0 = 2.31 \cdot 10^{-5}$ in both figures.

A quantity of primary interest is the renormalized temperature dependent tunneling rate, which can be experimentally determined by performing real-time measurements [7]. By dimensional analysis, for an ohmic heat bath it is given by $R(T) \sim \Delta(\omega = T)^2/T$ [1]. We computed it by integrating Eq. (5) (Fig. 2, bottom). As the most striking consequence of the Kondo effect, the logarithmic slope of $R(T)$ changes at $T \approx T_K$.

The energy scale Δ^* is determined [1] by the condition $\Delta(\omega = \Delta^*) \approx \Delta^*$, leading to

$$\Delta^* = \Delta_0 \left(C \frac{\Delta_0}{T_K} \right)^{\alpha_</(1-\alpha_<)}, \quad (8)$$

with C a constant of the order of unity. The constant $\alpha_<$ also characterizes the dissipative non-equilibrium dynamics of the TLS below T_K [1]. In Fig. 3 we show the rescaled spectral function, $\Delta^{*2} \varrho_T^z(\omega)/\omega$, related to the real part of the retarded response function. Without dissipation, $\alpha_< = 0$ ($F = 1$), the tunneling of the TLS is entirely coherent: the TLS oscillates between the two positions without damping, and the spectral function consists of two Dirac delta's. For $F < 1$ the coherence peak broadens: The oscillations become exponentially damped and at very long time scales (at $T = 0$) the correlation function behaves as $\langle T^z(t)T^z(0) \rangle \sim 1/t^2$. For even

smaller values of overlap the peak becomes completely invisible, and the formation of the Kondo compensation cloud suppresses the coherent motion.




FIG. 3. Rescaled spectral functions and the evolution of the coherence peak with increasing overlap. We computed Δ^* from Eq. (8) with $C = 0.0916$ and $T_K = e^{-1/2g}$.

In the general case, the TLS model is more complex than the one we discussed until now. The TLS is not symmetrical: $\Delta_z \neq 0$, and the couplings $J_{\pm}$ in Eq. (2) are not equal. The difference between $J_{\pm}$ leads to *two* subsequent Kondo effects at T_K 's associated with the two positions, and consequently changes in the logarithmic slope of $R(T)$ twice, while a finite Δ_z generates a new energy scale, below which the TLS freezes into one of the states $T_z = \pm 1/2$ [16].

In Eq. (2) we only took into account the exchange interaction. In reality, the TLS and the electrons interact through a local potential scattering as well. However, for a Kondo impurity this potential scattering is relatively small (of the order of the exchange coupling) in the exchange scattering-channel and since it remains essentially unrenormalized, its effect can be neglected compared to that of the exchange interaction. Potential scattering in *other* scattering channels may, however, be still present and shift α by a temperature-independent value $\alpha(T) \rightarrow \alpha(T) + \alpha_>$. Therefore, for small overlaps one can easily be in a situation, where the Kondo effect drives the TLS from a coherent ground state to an incoherent state with $\alpha > 1/2$ (see Fig. 2).

For small enough Kondo temperatures, the value of $\delta\sigma$ in Eq. (7) can be changed by an external magnetic field [23], which gradually destroys the Kondo effect, thereby changes the logarithmic slope of $R(T)$ to its value above T_K , and restores the coherent TLS motion [16].

In the experimental realizations mentioned in the introduction it is the Kondo spin that moves and carries the compensation cloud. Our discussion can be, however, easily generalized to the case of a TLS that happens to be close to a Kondo impurity [24,25]. In this case Friedel oscillations generated by the TLS modify the local density of states at the Kondo impurity and thus the exchange coupling will depend on the position of the TLS, leading to a Hamiltonian similar to Eq. (2).

It is also an interesting question whether the two-channel Kondo behavior found can possibly be observed

in realistic systems. Clearly, one needs an almost perfectly symmetrical TLS and a very small overlap F . In most cases F is typically large, excepting the magnetic impurity tunneling between an STM tip and a surface, where the electronic states coupled to the spin hardly overlap at the two positions and $F \approx 0$. In order to observe the two-channel Kondo behavior, one has to minimize the TLS asymmetry too: The asymmetry Δ_z can be tuned to zero with the external voltage, while $J_+ \approx J_-$ can be assured by using sufficiently rounded tip that is made of the same metal as the surface.

Acknowledgements: We are grateful to T. Costi, D.L. Cox, K. Damle, and E. Demler, for stimulating discussions, and especially J.P. Sethna and B.I. Halperin, for making useful suggestions. This research has been supported by NSF Grants Nos. DMR-9985978 and DMR97-14725, and Hungarian Grants No. OTKA F030041, T029813, and T29236.

-
- [1] A. J. Leggett et al., Rev. Mod. Phys. **59**, 1 (1987).
 - [2] Yu. Kagan and N. V. Prokof'ev, in *Quantum Tunneling in Condensed Media*, edited by Yu. Kagan and A. J. Leggett (North Holland, Amsterdam, 1992).
 - [3] Ulrich Weiss, *Quantum Dissipative Systems* (World Scientific, Singapore, 1993).
 - [4] F. Lesage et al., Phys. Rev. Lett. **76**, 3388 (1996).
 - [5] T. Costi and G. Zaránd, Phys. Rev. B **59**, 12398 (1999).
 - [6] N. Giordano in *Quantum Tunneling in Condensed Media* edited by Yu. Kagan and A. J. Leggett (North Holland, Amsterdam, 1992).
 - [7] K.S. Ralls and R. A. Buhrman, Phys. Rev. Lett. **60**, 998 (1988); B. Golding, et al., **68**, 998 (1992); D. H. Cobden et al., Phys. Rev. Lett. **71**, 4230 (1993).
 - [8] P.W. Anderson, Phys. Rev. Lett. **18**, 1049 (1967).
 - [9] D.L. Cox and A. Zawadowski, Adv. Phys. **47**, 599 (1998).
 - [10] A. Moustakas and D. S. Fisher, Phys. Rev. B **53**, 4300 (1996).
 - [11] G. Zaránd, Phys. Rev. B **52**, 13459 (1995).
 - [12] J. Ye, Phys. Rev. B **56**, 1316 (1997).
 - [13] A.A. Louis and J.P. Sethna, Phys. Rev. Lett. **74**, 1363 (1995).
 - [14] D.C. Ralph et al., Phys. Rev. Lett. **72**, 1064 (1994).
 - [15] A.J. Heeger et al., Rev. Mod. Phys. **60**, 781 (1988).
 - [16] L. Borda et al., (under preparation).
 - [17] W. Izumida and O. Sakai, Phys. Rev. B **62**, 10620 (2000).
 - [18] V. L. Libero and L. N. Oliveira, Phys. Rev. Lett. **65**, 2042 (1990).
 - [19] K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975); T. A. Costi in *Density Matrix Renormalization* edited by I. Peschel et al. (Springer 1999).
 - [20] B. A. Jones et al., Phys. Rev. Lett. **61**, 125 (1988).
 - [21] J. L. Cardy, *Scaling and Renormalization in Statistical Physics* (Cambridge Lecture Notes in Physics, 5).
 - [22] M. Yoshida, Phys. Rev. B **41**, 9403 (1990).
 - [23] P. Nozières, J. Low Temp. Phys. **17**, 31 (1974).
 - [24] R. J. P. Keijzers et al. Phys. Rev. Lett. **77**, 3411 (1996).
 - [25] G. Zaránd et al., Phys. Rev. Lett. **80**, 1353 (1998).